

TOPIC: CHI-SQUARE GOODNESS OF FIT TEST



Goodness of Fit Test

- ◆ Use a **Goodness of Fit test** to determine if _____ frequencies match "claimed" frequencies in a distribution.
 - Instead of testing just **one** parameter, you'll test the frequencies for _____ categories.

EXAMPLE

You roll a 6-sided die 60 times and list the observed frequencies in the table. Determine if this die is fair by testing the goodness of fit of the die rolls with a uniform distribution. Use $\alpha = 0.05$.

Roll	1	2	3	4	5	6
Frequency	13	12	1	11	14	9

Recall	Hypoth. Tests	New	Goodness of Fit Tests
1) Hypothesis	$H_0: \mu = \#$ $H_a: \mu \neq \#$	H_0 : observed frequencies match claimed distribution H_a : observed frequencies DO NOT match claimed distribution, i.e. at least _____ of the probabilities is different from claimed.	
2) Test Stat	$z = \frac{\bar{x} - \mu}{\sigma/\sqrt{n}}$	<p><i>If claimed prob's same:</i></p> <div style="display: flex; justify-content: space-around;"> <div> $\chi^2 = \sum \frac{(O - E)^2}{E}$ </div> <div> $E = \frac{n}{k}$ </div> <div> $O = \text{observed freq.}$ $E = \text{expected freq.}$ $n = \text{total sample size}$ $k = \# \text{ of categories}$ </div> </div> <p> $n = \underline{\hspace{2cm}}$ $k = \underline{\hspace{2cm}}$ $E = \underline{\hspace{2cm}}$ </p> $\chi^2 = \frac{(\underline{\hspace{1cm}} - \underline{\hspace{1cm}})^2}{10} + \frac{(\underline{\hspace{1cm}} - 10)^2}{10} + \frac{(\underline{\hspace{1cm}} - 10)^2}{10} + \frac{(\underline{\hspace{1cm}} - 10)^2}{10} + \frac{(\underline{\hspace{1cm}} - 10)^2}{10} + \frac{(\underline{\hspace{1cm}} - 10)^2}{10}$ $= \frac{\hspace{1cm}}{10} + \frac{\hspace{1cm}}{10} + \frac{\hspace{1cm}}{10} + \frac{\hspace{1cm}}{10} + \frac{\hspace{1cm}}{10} + \frac{\hspace{1cm}}{10}$	
3) P-value		<div style="display: flex; align-items: center;"> <div style="margin-right: 20px;"> $df = k - 1$ $df = \underline{\hspace{2cm}}$ </div> <div style="text-align: center;">  </div> <div> $P\text{-value} = \text{Area "beyond" } \chi^2$ $P\text{-value} = \underline{\hspace{2cm}}$ </div> </div>	
4) Conclusion	Because $P\text{-value} \dots$	<p>Because $P\text{-value}$ [< >] α, we [REJECT FAIL TO REJECT] H_0.</p> <p>There is [ENOUGH NOT ENOUGH] evidence to conclude that the observed frequencies do not match the claimed distribution, therefore it [IS IS NOT] a good fit and the die _____ fair.</p>	
Criteria	Random samples? <input type="checkbox"/> X is normal <input type="checkbox"/> OR $n > 30$?	Random Samples? <input type="checkbox"/> Observed freq. for each category? <input type="checkbox"/> $E \geq 5$ for each category? <input type="checkbox"/>	

TOPIC: CHI-SQUARE GOODNESS OF FIT TEST

PRACTICE

A gym owner wants to know if the gym has similar numbers of members across different age groups. The table shows the distribution of ages for members from a random survey. Write the null & alt. hypotheses to test the claim that the gym has equal numbers of members across all groups.

Age Group	18 – 25	26 – 35	36 – 45	46 – 55	56+
# of Members	54	46	53	49	48

H_0 :

H_a :

Find the χ^2 statistic to test the claim that the gym has equal numbers of members of all age ranges.

$n = \underline{\hspace{2cm}}$ $k = \underline{\hspace{2cm}}$

$E = \underline{\hspace{2cm}}$ $\chi^2 = \underline{\hspace{2cm}}$

Recall

$$E = \frac{n}{k}$$

(Expected Freq. – Uniform)

Recall

$$\chi^2 = \sum \frac{(O - E)^2}{E}$$

Using $\chi^2 = 0.92$ & $\alpha = 0.05$, test the claim that the gym has equal numbers of members of all age ranges.

$df = \underline{\hspace{2cm}}$

$P\text{-value} = \underline{\hspace{2cm}}$

Recall

$$df = k - 1$$

Because $P\text{-value}$ [< | >] α , we [**REJECT** | **FAIL TO REJECT**] H_0 .

There is [**ENOUGH** | **NOT ENOUGH**] evidence to conclude that the # of members is significantly different for at least one of the age groups at this gym.

So the claimed dist. [**IS** | **IS NOT**] a good fit.

Does this data set fit the criteria for a G.O.F. test?

Random Samples? ☐

Observed freq. for each category? ☐

$E \geq 5$ for each category? ☐

TOPIC: CHI-SQUARE GOODNESS OF FIT TEST

EXAMPLE

A company runs a customer satisfaction survey where customers rate their experience. The manager claims that responses will NOT be the same across all 5 rating categories. A random sample of 100 survey responses is collected with the following observed frequencies. Using $\alpha = 0.05$, test the manager's claim.

Survey Response	Very Poor	Poor	Neutral	Good	Very Good
Frequency	13	14	26	29	18

Random Samples? ☐

Observed freq. for each category? ☐

$E \geq 5$ for each category? ☐

H_0 :

H_a :

$n = \rule{1cm}{0.4pt}$ $k = \rule{1cm}{0.4pt}$ $E = \rule{1cm}{0.4pt}$

Recall

$$\chi^2 = \sum \frac{(O - E)^2}{E}$$
$$df = k - 1$$

Recall

$$E = \frac{n}{k}$$

(Expected Freq. – Uniform)

Because P -value [< | >] α , we [REJECT | FAIL TO REJECT] H_0 .
There is [ENOUGH | NOT ENOUGH] evidence of H_a .
So the claimed dist. [IS | IS NOT] a good fit.

TOPIC: CHI-SQUARE GOODNESS OF FIT TEST

Goodness of Fit Test: Unequal Probabilities

- ◆ If claimed probabilities AREN'T equal, find expected freq's using *given probabilities* instead of # of categories.

EXAMPLE

According to Benford's Law, the 1st digits of large numbers in real-world data sets follow the probability distribution in the table. Also listed in the table are the 1st-digit frequencies of the populations of 100 random cities across Europe. Find the test statistic that would be used to test if these digits follow Benford's Law.

1 st Digit	1	2	3	4	5	6	7	8	9
Frequency (<i>O</i>)	18	14	11	13	12	9	5	8	10
Benford's Law Prob's (<i>p</i>)	.301	.176	.125	.097	.079	.067	.058	.051	.046
Exp. Freq. (<i>E</i>)									
$\frac{(O - E)^2}{E}$									

New
 $E = np$ $n = \underline{\hspace{2cm}}$

Recall
$$\chi^2 = \sum \frac{(O - E)^2}{E}$$

$\chi^2 = \underline{\hspace{2cm}}$

PRACTICE

A marketing associate for a supermarket chain wants to determine how many of each snack type to cook. According to previous market research, customers' preferences tend to follow the distribution in the table. If approximately 200 snack items are purchased in a day, what is the expected frequency of each snack type?

1 st Digit	Chips	Cookies	Crackers	Nuts	Granola Bars
Preferences from Research	36%	21%	12%	8%	23%
Expected Sales					

Recall
 $E = \frac{n}{k}$ (claimed prob's SAME)
 $E = np$ (claimed prob's DIFF)